

# PIGEON

## MEDICINES AND SUPPLEMENTS





## 01. Therapeutics

### Antibiotics for internal administration

Antibiotic powders for oral solutions .....	02
Antibiotic solutions .....	20
Antibiotic tablets .....	21

### Antibiotics for external administration

Antibiotic cutaneous powders.....	30
Antibiotic ointments and emulsions.....	31
Antibiotic sprays and solutions.....	33
Antibiotic ear drops.....	35

### Non-antibiotics for external administration

Cutaneous ointments.....	36
Cutaneous sprays and solutions.....	38

## 02. Nutritionals

Tablets.....	40
Solutions and powders for oral administration .....	41

## 03. Parasitics

Cutaneous powders.....	44
Cutaneous sprays and solutions.....	45
Powders, tablets and solutions for oral administration .....	53
Liquid for water solution for spraying the premises .....	56





Package size:

10g 20g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Colitetraviv

**Antibiotics & Vitamins complex for treatment of colibacteriosis, salmonellosis, pasteurellosis, mycoplasmosis. Power for oral administration with drinking water.**

**Composition in 100g:** Oxytetracycline hydrochloride: 5.0g  
Colistinsulphate: 3 500 000 UI, Vitamin A: 50 000 UI, Vitamin D3: 5 000 UI, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons, decorative birds, decorative rabbits, guinea pigs.

**Indications:** Colitetraviv is a combination of Oxytetracycline hydrochloride, Colistin sulphate and balanced vitamin complex. Oxytetracycline hydrochloride is a broad-spectrum antibiotic, and the synergism between the two antibiotics enhances the effect of the treatment. It is highly effective against colibacteriosis; salmonellosis; pasteurellosis; mycoplasmosis.

**Contraindications:** Do not combine with products containing salicylic acid.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/1L of water for 4 to 6 consecutive days. Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Metronidazol

**Against histomonas, trichomonas, lamblia, amoebae. Powder for oral administration with the drinking water.**

**Composition in 100g:** Metronidazole 10g.

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** Metronidazole has an antiprotozoal action against histomonas, trichomonas, lamblia, amoebae; has systematic antiprotozoal effect. Its toxicity for birds is low.

Do not administer to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** For treatment – 0.6g/1L water for 9 consecutive days, then in prophylactic doses (0.2g/L water) for 5 days.

For prophylaxis - 0.2g/1L of water for 8 weeks.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Metronidazol Forte

**Against histomonas, trichomonas, lamblia, amoebae. Powder for oral administration with the drinking water.**

**Composition in 100g:** Metronidazole 98%

**Excipients:** None

**Target species:** Pigeons.

**Indications:** Metronidazole has an antiprotozoal action against histomonas, trichomonas, lamblia, amoebae; has systematic antiprotozoal effect. Its toxicity for birds is low.

Do not administer to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water

**Dosage:** For treatment – 1g/5L water for 3-5 consecutive days.

For prophylaxis – 1g/10L water for 3-5 consecutive days.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 100g 1Kg

Store below 28°C in cool dry place

## Ampicillin 50%

**Antibiotic for treatment of bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Ampicillin trihydrate 4 000 000 IU

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** For treatment of diseases caused by ampicillin-sensitive microorganisms like bronchopneumonias; gastrointestinal diseases caused by Salmonella and E. Coli; against pasteurellosis and spirochaetosis.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** For treatment – 0.6g/1L water for 9 consecutive days, then in prophylactic doses (0.2g/L water) for 5 days. For prophylaxis - 0.2g/1L of water for 8 weeks. Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Oxytetracycline HCL

Antibiotic for treatment and prophylaxis of primary and secondary bacterial infections. Powder for oral administration with the drinking water.

**Composition in 100g:** Oxytetracycline hydrochloride: 95%

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** For treatment and prophylaxis of infectious diseases of the gastrointestinal, respiratory and urogenital tracts, arthritis and secondary bacterial infections due to viral diseases.

**Mechanism of action:** Oxytetracycline hydrochloride is a broad-spectrum antibiotic that inhibits the development and reproduction of a large number of microorganisms (Gram-positive and some Gram-negative bacteria, mycoplasmas, rickettsiae, some large viruses and protozoans).

**Route of administration:** Internally, dissolved in the drinking water

**Dosage:** 2.5-4g/10L of water for 5 consecutive days.

Solution should be prepared daily before use.

If condition does not improve within three days, an antibiogram should be made and treatment should be changed if necessary.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 100g 100g 250g 500g 1Kg

Store below 28°C in cool dry place

## Oxytetracycline 50% + Vitamins

Antibiotic & Vitamin complex for treatment and prophylaxis of infectious diseases. Powder for oral administration with the drinking water.

**Composition in 100g:** Oxytetracycline hydrochloride: 50g, Vitamin A: 75 000 IU, Vitamin D3: 7500 IU, Vitamin E acetate: 12.5mg, Vitamin K: 7.5mg, Vitamin PP: 1mg, Vitamin B1: 10mg, Vitamin B2: 20mg, Vitamin B6: 5mg, Vitamin B12: 0.5mg, Vitamin C: 125mg

**Excipients:** Dextrose monohydrate to 100g

**Target species:** Pigeons.

**Indications:** For treatment and prophylaxis of infectious diseases of the gastrointestinal, respiratory and urogenital tracts, arthritis and secondary bacterial infections due to viral diseases. The vitamin complex increases resistance during and after diseases, improves the metabolism and appetite, reduces the apathy and fatigue.

**Dosage:** For treatment: 1g/2L water for 5 consecutive days

For prophylaxis: 1g/4L of water for 5 consecutive days

Solution should be prepared daily before use.

If condition does not improve within three days, an antibiogram should be made and treatment should be changed if necessary.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## OxytetraVIT

Antibiotic & Vitamin complex for treatment and prophylaxis of infectious diseases. Powder for oral administration with the drinking water.

**Composition in 100g:** Oxytetracycline hydrochloride: 5.0g, Vitamin A: 50000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg.

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** OxytetraVIT is a combination of Oxytetracycline hydrochloride and balanced vitamin complex for treatment and prophylaxis of bacterial infections sensitive to oxytetracycline like infectious synovitis caused by Mycoplasma synoviae; chronic respiratory disease (CRD) and infectious aerosacculitis caused by M.gallisepticum and E. Coli.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/2L of water for 4 to 6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Gentamicin Pulvis

Antibiotic for treatment of bacterial infections  
Powder for oral administration with the drinking water

**Composition in 100g:** Gentamicin sulfate: 1 600 000 IU.

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** Gentamicin is a broad spectrum antibiotic with a bactericidal activity against Gram-positive and Gram-negative microorganisms like staphylococci, streptococci, colibacteria, proteus, brucellas, salmonella, etc., including antibiotic-resistant microorganisms. For treatment of digestive, urinary and pulmonary infections, respiratory mycoplasmosis, sepsis.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 3-5g/3L of water for 5 consecutive days. Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Gentavit

**Antibiotic & Vitamin complex for treatment and prophylaxis of infectious diseases. Powder for oral administration with the drinking water.**

**Composition in 100g:** Gentamicin sulfate: 600 000 IU, Vitamin A: 75 000 IU, Vitamin D3: 7 500 IU, Vitamin E acetate: 10mg, Vitamin K3: 15mg, Vitamin PP: 190mg, Vitamin B1: 10mg, Vitamin B2: 37mg, D-Pantothenic Acid: 38mg, Vitamin B6: 15mg, Vitamin B12: 0,04mg, Vitamin C: 375mg, Folic acid: 1.5mg, Vitamin H: 0.5mg

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** Gentavit is a broad-spectrum antibiotic with bactericidal action against Gram-positive and Gram-negative microorganisms like staphylococci, streptococci, colibacteria, proteus, brucellas, salmonella, etc., including against microorganisms resistant to other antibiotics. For treatment of digestive, urinary and pulmonary infections, respiratory mycoplasmosis, sepsis. The vitamins complex increases resistance during and after illnesses, improves metabolism and appetite, reduces apathy and fatigue.

**Route of administration:** Internally, dissolved in the drinking water

**Dosage:** 3-5g/3L of water for 5 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Tetraneovit

**Antibiotics & Vitamin complex for treatment and prophylaxis of infectious diseases Powder for oral administration with the drinking water.**

**Composition in 100g:** Oxytetracycline hydrochloride: 6.00g, Neomycin sulphate: 0.500g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** For treatment of chronic respiratory diseases, synovitis, sinusitis and other infections caused by microorganisms sensitive to oxytetracycline and neomycin.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/1L of water for 4-6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 1Kg

Store below 28°C in cool dry place

## Tylofarm

**Antibiotic for treatment of bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Tylosin tartrate: 50g.

**Excipients:** Dextrose monohydrate to 10g.

**Target species:** Pigeons.

**Indications:** Tylosin is effective against Gram-positive and Gram-negative microorganisms, protozoa and large viruses, clostridia, brucellas, rickettsiae, hemophiles, spirochetes, chlamydia from the psittacoses group and others. The performance of tylosin against mycoplasmas is very good, better than the performance of many chemotherapeutic agents. When administered internally it is well absorbed so bacteriostatic concentrations in blood and tissues are reached. Tylosin can be combined with sulfonamides, tetracycline, nitrofurantoin, erythromycin, spectinomycin. Solution is thermostable but sensitive to light.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/1L of water for 4-6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Tylocyclin + Vitamins

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Tylosin tartrate: 20g, Chlortetracycline hydrochloride: 10g, Vitamin A: 75 000 IU, Vitamin D3: 7500 IU, Vitamin E acetate: 2.5mg, Vitamin K: 7.5mg, Vitamin PP: 1mg, Vitamin B1: 10mg, Vitamin B2: 20mg, Vitamin B6: 5mg, Vitamin B12: 0.5mg, Vitamin C: 125mg

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** A combination of two antibiotics with a broad spectrum of action for treatment of primary and secondary bacterial infections of the respiratory and gastrointestinal tracts. The vitamin complex increases resistance during and after illnesses, improves metabolism and appetite, reduces apathy and fatigue.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/1L of water for 5 consecutive days.

**Withdrawal period:** Not applicable.





Package size:

10g 100g 200g 1Kg

Store below 28°C in cool dry place

## Colistin

**Antibiotic for treatment and prophylaxis of bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Colistin sulphate 750 000 UI

**Target species:** Pigeons.

**Indications:** In cases of intestinal infections caused by Gram-negative and colistin-sensitive bacteria.

**Contraindications:** Colistin should not be administered in cases of proven polymyxin resistance and renal insufficiency.

The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** For treatment: 1-2g/1L of water for 3-5 consecutive days. For prophylaxis: 0.5-1g/1L of water for 3-5 consecutive days. Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 1Kg

Store below 28°C in cool dry place

## Vitafur

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Neomycin sulfate: 6g, Oxytetracycline Hydrochloride: 3g, Furaltadon Hydrochloride: 2g, Vitamin A: 50000 IU, Vitamin D3: 5000 IU, Vitamin E acetate: 7mg, Vitamin K3: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, D-Pantothenic Acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg.

**Excipients:** Dextrose monohydrate to 10g.

**Target species:** Pigeons and decorative birds.

**Indications:** Oxytetracycline is active against Gram-positive and Gram-negative bacteria. It acts against E-coli, salmonella, staphylococcus, streptococcus. Furaltadon is a chemotherapeutic agent with bacteriostatic action against salmonella and E-coli. For treatment of intestinal, respiratory and urogenital infections caused by bacteria sensitive to oxytetracycline, neomycin and furaltadon.

**Route of administration:** Internally, with the feed or with the drinking water.

**Dosage:** Young birds – 2-6g/1kg fodder adult birds 1-2g/1L of water for 5-6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Enterogard Farma

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Metronidazole: 10g, Oxytetracycline hydrochloride: 6g, Furaltadon hydrochloride: 2g, Vitamin A: 50000 IU, Vitamin D3: 5000 IU, Vitamin E acetate: 7mg, Vitamin K3: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, D-Pantothenic Acid: 50mg, Nicotinic acid: 130mg, Biotin 0.35mg

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** Oxytetracycline is active mainly against Gram-positive and some Gram-negative bacteria, including some anaerobes; E. coli, Salmonella spp., Staphylococcus spp., Streptococcus spp.

Furaltadon is active against E. coli, Salmonella spp.

Metronidazole has an antiprotozoal effect against histomonas, trichomonas, lamblia and amoebas. ENTEROGARD FARMA acts on various microorganisms that cause infections on the gastrointestinal mucosa and intestines.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1-2g/1L water for 5-6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Tiamol 45

**Antibiotic for treatment and prophylaxis of bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Tiamulin Hydrogen Fumarate: 45g.

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of diseases of the respiratory tract caused by mycoplasmas, E-coli, Pasteurella and Haemophilus gallinarum.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** For treatment: 5g/10L of water for 3-5 consecutive days.

For prophylaxis: 2.5g/5L of water for 3-5 consecutive days.

Recommendation: first add a small amount of warm water to the powder, stir well and then add the rest of the water.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 1Kg

Store below 28°C in cool dry place

## Tyldoxin – Vita

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Doxycycline Hyclate: 5g, Tylosin tartarate: 20g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.0025mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg

**Excipient:** Dextrose monohydrate to 100g

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of primary and secondary bacterial infections of the respiratory and gastrointestinal tracts, mycoplasmosis, clambia, ornithosis. The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 5g/2L water for 3-7 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Furaltadon 6

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Furaltadone HCl: 6g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E acetate: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg.

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** Furaltadone 6 is one of the best products for treatment of bacterial infections like salmonella and E-coli, easy to use, very effective and affordable.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 5g/1L of water for 3 to 7 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g

Store below 28°C in cool dry place

## Ornivit

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Doxycycline Hyclate: 6g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** Doxycycline is absorbed quickly and completely, effectively penetrates into the lung tissue which makes it effective for lung infections. It is used for ornithosis, mycoplasmosis, and breathing problems.

The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 5g/1L of water for 3-7 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

20g 100g 1Kg

Store below 28°C in cool dry place

## Furaltadon 100 %

**Antibiotic for treatment of bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Furaltadone HCl: 99.9%

**Target species:** Pigeons and decorative birds.

**Indications:** Furaltadone 100% is one of best products for treatment of bacterial infections like salmonella and E-coli, easy to use, very effective and affordable.

The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 5g/4-5L of water for 3 to 7 consecutive days.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 100g 1Kg

Store below 28°C in cool dry place

## Doxymulin

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Doxycycline Hydrate: 5g, Tiamulin hydrogen fumarate: 5g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0,35mg.

**Excipient:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment of primary and secondary bacterial infections of the respiratory and gastrointestinal tracts caused by gram-positive and gram-negative microorganisms.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1-2g/1L of drinking water.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Chloramphenicol + Vitamins

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Chloramphenicol: 5.0g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Vitamin C: 250mg, Folic acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg .

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment of bacterial infections caused by chloramphenicol sensitive microorganisms such as salmonellosis, respiratory and gastrointestinal tract infections, contagious runny nose, and septicemia.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/1L of water for 4 to 6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 100g 500g 100g

Store below 28°C in cool dry place

## Doxycyclin 20%

**Antibiotic for treatment of infections of the respiratory and gastrointestinal tracts. Powder for oral administration with the drinking water**

**Composition in 100g:** Doxycycline hyclate 20g

**Excipients:** Lactose to 100g

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment of infections of the respiratory and gastrointestinal tracts caused by microorganisms sensitive to doxycycline.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 0.5-1g/1L of drinking water for 3-5 consecutive days. Solution should be prepared daily before use

**Withdrawal period:** Not applicable.



Package size:

20g 100g 1Kg

Store below 28°C in cool dry place

## Chloramphenicol 100%

**Antibiotic for treatment of bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Chloramphenicol: 98.0g

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment of bacterial infections caused by chloramphenicol sensitive microorganisms such as salmonellosis, respiratory and gastrointestinal tract infections, contagious runny nose, and septicemia.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/4-5L of water for 4 to 6 consecutive days.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.





Package size:

10g 20g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Paramixovit

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Furaltadone HCl: 10g, Chloramphenicol: 7g, Colistin sulphate: 5g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin K: 10mg, Vitamin C: 250mg, Vitamin B1: 6,5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic Acid: 1mg, Pantothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg

**Excipient:** Dextrose monohydrate to 100g

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment of Salmonellosis (paratyphoid), E-coli, bacterial infections of the gastrointestinal tract.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1 - 2g/1L of water for 4 to 6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Avigard + Vitamins

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Neomycin sulfate: 10g, Tylosin tartarat: 7.0g, Erythromycin thiocyanate: 2.0g, Vitamin A: 75 000 IU, Vitamin D3: 7 500 IU, Vitamin E: 12.5mg, Vitamin K: 7.5mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 1mg, Vitamin PP: 1mg.

**Excipients:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** For treatment and prophylaxis of mycoplasmosis (chronic respiratory disease), salmonellosis, pasteurellosis, colibacteriosis, secondary microbial infections.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1-2g/1L of water for 5-7 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 500g 1Kg

Store below 28°C in cool dry place

## Respigard + Vitamins

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Tetracycline hydrochloride: 10.0g, Tylosin tartarat: 7.0g, Erythromycin thiocyanate: 2.0g, Vitamin A : 75 000 IU, Vitamin D3: 7 500 IU, Vitamin E: 12,5mg, Vitamin K: 7.5mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 1mg, Vitamin PP: 1mg

**Excipients:** Dextrose monohydrate to 100g

**Target species:** Pigeons

**Indications:** For treatment and prophylaxis of respiratory organs, mycoplasmosis (chronic respiratory disease - CRD), ornithosis, secondary microbial infection.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1 - 2g /1L of water for 5 - 7 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

20g 100g 1Kg

Store below 28°C in cool dry place

## Ronidazol 100%

**Antibiotic for treatment and prophylaxis of trichomoniasis and hexamitiasis. Powder for oral administration with the drinking water.**

**Composition in 100g:** Ronidazole: 10g.

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of trichomoniasis and hexamitiasis or hexamitosis (gangrene).

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 2g/4-5L water for 3 to 5 consecutive days.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 1Kg

Store below 28°C in cool dry place

## Ronidazol 10%

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water**

**Composition in 100g:** Ronidazole: 10g

**Excipient:** Dextrose monohydrate to 100g

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of trichomoniasis and hexamitiasis or hexamitosis (gangrene).

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 2g/1L water for 3 to 5 consecutive days.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

20g 50g 100g 500g

Store below 28°C in cool dry place

## Sulfaguanidin

**Antibiotic for treatment of gastrointestinal diseases with infectious nature. Powder for oral administration with the feed.**

**Composition in 100g:** p-Aminobenzol-sulfonylguanidinum monohydratum.

**Target species:** Pigeons.

**Indications:** For treatment of gastrointestinal diseases with infectious nature. The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, with the feed.

**Dosage:** 10g/20 kg feed /0.5% in the fodder/ for 4 to 7 consecutive days.

**Precautions for use:** Wear personal protective equipment while handling the product!

**Withdrawal period:** Not applicable.



Package size:

10g 20g 100g 1Kg

Store below 28°C in cool dry place

## Ronifur

**Antibiotic for treatment and prophylaxis of bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Ronidazole: 5g, Furaltadone HCl: 5g

**Excipients:** Dextrose monohydrate to 100g

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of trichomoniasis, gangrene, E-coli, paratyphoid, coccidiosis, gastro-intestinal infections, diarrhea.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 5g (1 teaspoon)/4L water for 3 to 5 consecutive days.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 50g 100g 1Kg

Store below 28°C in cool dry place

## Erythrofurox + Vitamins

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Oxytetracycline hydrochloridum: 1.65g, Furaltadone hydrochloridum: 5.50g, Erythromycin: 3.70g, Vitamin A: 75 000 IU, Vitamin D3: 7 500 IU, Vitamin E: 12.5mg, Vitamin K: 7.5mg, Vitamin C: 250mg, Vitamin B1: 10mg, Vitamin B2: 20mg, Vitamin B6: 5mg, Vitamin B12: 0.5mg, Vitamin PP: 1mg.

**Excipient:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** For prophylaxis and treatment of respiratory infections (ornithosis), diseases caused by Salmonella and E. coli, and other bacterial infections. The preparation helps the general recovery from and after diseases, the vitamin complex increases resistance during and after illnesses, improves metabolism and appetite, reduces apathy and fatigue.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** For treatment: 5g (1 spoon)/2L of water for 5-7 consecutive days. Repeat if necessary. For prophylaxis: 5g (1 spoon)/2L of water for 1-2 consecutive days every two weeks.

**Withdrawal period:** Not applicable.





Package size:

1g	6g	10g	15g	50g	100g
		250g	500g	1Kg	

Store below 28°C in cool dry place

## Streptomycin

**Antibiotic for treatment of bacterial infections. Powder for oral administration with the drinking water**

**Composition in 1mg:** Streptomycin sulfate: 720 UI/mg.

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment of fowlpox in pigeons, diarrhea of different nature, in chronic forms of Salmonella and salmonella arthritis as it acts primarily on Gram-negative but also on some Gram-positive bacteria and acid-resistant microorganisms..

In some cases besides dissolved in water, streptomycin should be administered intramuscularly as well.

Streptomycin can be combined with Penicillin, Neomycin, Bacitracin, Sulfonamides.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g/1L water for 3 to 5 consecutive days

**Withdrawal period:** Not applicable.



Package size:

6g	10g	20g	50g	100g
		250g	500g	1Kg

Store below 28°C in cool dry place

## Penicillin

**Antibiotic for treatment of streptococcosis**

**Powder for oral administration with the drinking water or the feed.**

**Composition in 1g:** Penicillin: 1 000 000 UI/g

**Target species:** Pigeons.

**Indications:** For treatment of streptococcosis in pigeons.

The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, with the feed.

**Dosage:** 5-6g/1kg of feed for 4 - 5 consecutive days.

OR 3-6mg per pigeon for 5 consecutive days into crumbs of bread.

OR 1-2g/1L of drinking water.

**Precautions for use:** Wear personal protective equipment while handling the product!

**Withdrawal period:** Not applicable.



Package size:

10g	50g	100g	1Kg
-----	-----	------	-----

Store below 28°C in cool dry place

## Amoxicillin

**Antibiotic for treatment of primary and secondary infections of the respiratory, gastrointestinal and urogenital tracts. Powder for oral administration with the drinking water.**

**Composition in 1g:** Amoxicillin trihydrate: 500mg/g

**Excipients:** Citric Acid Anhydrous to 1g

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment of primary and secondary infections of the respiratory, gastrointestinal and urogenital tracts caused by microorganisms sensitive to amoxicillin like salmonella, staphylococci, streptococci, E. coli and others.

**Contraindications:** Do not combine with tetracycline and macrolide antibiotics.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 3-5g /1L of water

**Withdrawal period:** Not applicable.



Package size:

10g	20g	50g	100g
-----	-----	-----	------

Store below 28°C in cool dry place

## Tricho-Cocci Vit

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Amprolium HCL 40mg, Ronidazole 30mg,

Vitamins – A, D<sub>3</sub>, E, B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, B<sub>12</sub>, K<sub>3</sub>, C, Nicotin amide, folic acid, biotin, methionine, lysine, sodium, potassium, magnesium, ferrous sulphate, D-calcium pantothenate.

**Excipient:** Dextrose monohydrate to 100g

**Target species:** Pigeons

**Indications:** For treatment of trichomonosis, coccidiosis, hexamitose and other protozoal infections caused by flagellates like trichomoniasis, hexamitose and coccidiosis. Helps improving digestion.

The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 5g/2L water for 5-7 consecutive days.

Solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Biostim

**Antibiotic & Vitamin complex for treatment of primary and secondary bacterial infections. Powder for oral administration with the drinking water.**

**Composition in 100g:** Chlortetracycline HCl: 5.5g, Vitamin A: 50 000 IU, Vitamin D3: 5 000 IU, Vitamin E: 7mg, Vitamin C: 250mg, Vitamin B1: 6.5mg, Vitamin B2: 25mg, Vitamin B6: 10mg, Vitamin B12: 0.025mg, Folic acid: 1mg, Panthothenic acid: 50mg, Nicotinic acid: 130mg, Biotin: 0.35mg

**Target species:** Pigeons

**Indications:** Broad-spectrum antibiotic with activity mainly against Gram-positive and some Gram-negative bacteria, including and some anaerobes. For treatment of ornithosis, gastrointestinal, respiratory and general bacterial infections. The vitamin complex boosts the metabolism and improves resistance to infections.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1-2g/1L water for 4 to 6 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Avigard

**Antibiotic for treatment and prophylaxis of primary and secondary microbial infections.**

**Composition for one tablet 150mg:** Neomycin sulfate: 10mg, Tylosin tartarat: 6.4mg, Erythromycin thiocyanate: 1.6mg, Vitamin A, Vitamin: D3, Vitamin: E, Vitamin: K, Vitamin: C, Vitamin: B1, Vitamin: B2, Vitamin: B6, Vitamin B12.

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of mycoplasmosis (chronic respiratory disease), salmonellosis, pasteurellosis, colibacteriosis, secondary microbial infections.

**Route of administration:** Orally.

**Dosage:** 1 tablet once a day for 3 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10ml 20ml 50ml 100ml 500ml 1L

Store below 28°C in cool dry place

## Enrofloxacin 10%

**For treatment of colibacillosis, salmonellosis, mycoplasmosis, chronic respiratory disease and other infections. Antibiotic oral solution.**

**Composition in 1ml:** Enrofloxacin: 100mg.

**Target species:** Pigeons.

**Indications:** For treatment of colibacillosis, salmonellosis, mycoplasmosis, chronic respiratory disease (CRD), staphylococcal infections.

**Contraindications:** Do not use together with chloramphenicol, tetracycline, macrolide antibiotics.

The product should not be administered to poultry intended for human consumption.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 0.5 ml / 1 L water for 3 to 5 days.

The solution should be prepared daily before use.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 130pcs

Store below 28°C in cool dry place

## Enterogard Farma

**Antibiotic tablets for treatment of bacterial infections.**

**Composition for one tablet 750mg:** Metronidazol: 75mg, Oxytetracycline hydrochloride: 37.5mg, Furaltadon hydrochloride: 15mg, Vitamin: A, Vitamin: D3, Vitamin: E, Vitamin: K, Vitamin: C, Vitamin: B1, Vitamin: B2, Vitamin: B6, Vitamin: B12, Folic acid, Nicotinic acid, Biotin

**Target species:** Pigeons, decorative birds and decorative animals (guinea pigs, hamsters, decorative rabbits).

**Indications:** Used for treatment of colibacteriosis; salmonellosis; pasteurellosis; mycoplasmosis, staphylococci, pasteurellosis, etc. Affects various microorganisms that cause infections of the gastrointestinal mucosa and intestines. Suitable for treatment of primary and secondary infections caused by microorganisms sensitive to the combination of metronidazol, oxytetracycline and furaltadon, like colibacteriosis; salmonellosis; pasteurellosis; mycoplasmosis, staphylococci, pasteurellosis, etc.

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablet once a day for 5-10 consecutive days. Guinea pigs, hamsters, decorative bunnies - 1 - 2 tablets once a day for 5-10 consecutive days.

**Withdrawal period:** Not applicable.





Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Vitafur

**Antibiotic tablets against intestinal, respiratory and urogenital diseases.**

**Composition for one tablet 750mg:** Neomycin sulfate: 44.7mg, Oxytetracycline Hydrochloride: 22.5mg, Furaltadon Hydrochloride: 19.2mg, Vitamin: A, Vitamin: D3, Vitamin: E, Vitamin: K, Vitamin: C, Vitamin: B1, Vitamin: B2, Vitamin: B6, Vitamin: B12, Folic acid, Nicotinic acid, Biotin.

**Target species:** Pigeons and decorative birds.

**Indications:** Suitable for treatment of intestinal, respiratory and urogenital diseases.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablets once a day for 5-10 consecutive days.

Guinea pigs, hamsters, decorative bunnies - 1 - 2 tablets once a day for 5-10 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 130pcs

Store below 28°C in cool dry place

## Metronidazol

**Antibiotic against trichomonosis.**

**Composition for one tablet 750mg:** Metronidazol: 75mg

**Target species:** Pigeons, decorative birds and decorative animals (guinea pigs, hamsters, decorative rabbits).

**Indications:** For treatment of trichomonosis.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablet once a day for 5-10 days.

Guinea pigs, hamsters, decorative rabbits - 1 - 2 tablets once a day for 5 - 10 days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Enrofloxacin

**Antibiotic tablets against bacterial infections.**

**Composition for one tablet 750mg:** Enrofloxacin: 15mg

**Target species:** Pigeons, decorative birds and decorative animals (guinea pigs, hamsters, decorative rabbits).

**Indications:** For treatment of colibacteriosis, salmonellosis, pasteurellosis, mycoplasmosis, staphylococci, pasteurellosis, etc.

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablet once a day for 5-10 days.

Guinea pigs, hamsters, decorative bunnies - 1 - 2 tablets once a day for 5-10 days.

**Precautions for use:** Wear personal protective equipment while handling the product!

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 130pcs

Store below 28°C in cool dry place

## Doxymulin

**Antibiotic against primary and secondary bacterial infections of the respiratory and gastrointestinal tract.**

**Composition for one tablet 750mg:** Doxycycline Hyclate: 75mg,

Tiamulin hydrogen fumarat: 75mg

**Target species:** Pigeons, decorative birds and decorative rabbits.

**Indications:** For treatment of primary and secondary bacterial infections of the respiratory and gastrointestinal tract caused by gram-positive and gram-negative microorganisms.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablet once a day for 5-10 days.

Decorative rabbits: 1 - 2 tablets once a day for 5-10 days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Sulfaguanidin

Antibiotic tablets for treatment of gastrointestinal diseases and coccidiosis

**Composition for one tablet 750mg:** Sulfaguanidin: 150mg

**Target species:** Pigeons, decorative birds and decorative rabbits.

**Indications:** For the treatment of gastrointestinal diseases and coccidiosis in exotic birds, pigeons and decorative rabbits.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablet once a day for 5 - 10 days.

Decorative rabbits: 1 - 2 tablets once a day for 5 - 10 days.

**Withdrawal period:** Not applicable.

## Tylocyclin

Antibiotic tablets against primary and secondary bacterial infections of the respiratory and gastrointestinal tract and ornithosis.

**Composition for one tablet 750mg:** Tylosin tartrate: 20mg,

Chlortetracycline hydrochloride: 10mg

**Target species:** Pigeons, decorative birds and decorative rabbits.

**Indications:** A combination of two antibiotics with a broad spectrum of action. For treatment of primary and secondary bacterial infections of the respiratory and gastrointestinal tract, ornithosis.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablets once a day for 5 - 10 days.

Decorative rabbits: 1 - 2 tablets once a day for 5 - 10 days.

**Precautions for use:** Wear personal protective equipment while handling the product!

**Withdrawal period:** Not applicable.



Package size:

50pcs 100pcs

Store below 28°C in cool dry place

## Furaltadon

Antibiotic tablets against primary and secondary bacterial infections of the respiratory and gastrointestinal tract.

**Composition for one tablet 750mg:** Furaltadone HCl: 50mg.

**Target species:** Pigeons, decorative birds and decorative rabbits.

**Indications:** For treatment of primary and secondary bacterial infections of the respiratory and gastrointestinal tract.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablet once a day for 5 - 10 days.

Decorative rabbits: 1 - 2 tablets once a day for 5-10 days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 130pcs

Store below 28°C in cool dry place

## Tylodoxin

Antibiotic tablets for treatment and prophylaxis of primary and secondary bacterial infections of the respiratory and gastrointestinal tract.

**Composition for one tablet 150mg:** Doxycycline Hyclate: 5mg, Tylosin tartrate: 20mg.

**Target species:** Pigeons, decorative birds and decorative rabbits.

**Indications:** An extremely effective combination for treatment and prophylaxis of mycoplasmosis, clambia, ornithosis and other primary and secondary bacterial infections of the respiratory and gastrointestinal tract.

*The product should not be administered to poultry intended for human consumption*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: ½ - 1 tablet once a day for 5-10 days.

Guinea pigs, hamsters, decorative bunnies - 1 - 2 tablets once a day for 5-10 days.

**Withdrawal period:** Not applicable.



Package size:

50pcs 100pcs

Store below 28°C in cool dry place





Package size:

20pcs 30pcs 130pcs

Store below 28°C in cool dry place

## Paramixovit

Antibiotic tablets and vitamin complex for treatment and prophylaxis of primary and secondary bacterial infections of the gastrointestinal tract.

**Composition for one tablet 150mg:** Furaltadone HCl, Chloramphenicol, Colistin sulphate, Complex of vitamins.

**Target species:** Pigeons.

**Indications:** For treatment and prophylaxis of salmonellosis, E-coli, severe diarrhea, intestinal infections caused by bacteria.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** 1 tablet once a day for 3-5 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

50pcs 100pcs

Store below 28°C in cool dry place

## Enrofloxacin

Antibiotic tablets for prophylaxis and treatment of primary and secondary bacterial infections of the respiratory and gastrointestinal tract.

**Composition for one tablet 150mg:** Enrofloxacin: 15mg, Colistin sulphate: 15mg, Vitamin complex.

**Target species:** Pigeons, decorative birds and decorative rabbits.

**Indications:** Effective against respiratory and gastro-intestinal infections like salmonellosis, mycoplasmosis, cholera, staphylococci, streptococci, infectious rhinitis.

**Route of administration:** Orally.

**Dosage:** For prophylaxis: 1 - 2 tablets once a day for 3 consecutive days.

For treatment- 1 - 2 tablets once a day for 5-10 consecutive days

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 50pcs 100pcs

Store below 28°C in cool dry place

## Ronidazol

Antibiotic tablets against trichomoniasis and hexamitiasis.

**Composition for one tablet 150mg:** Ronidazole: 15mg

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of trichomoniasis and hexamitiasis or hexamitosis (gangrene).

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** 1-2 tablets once a day for 5-10 days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 50pcs 100pcs

Store below 28°C in cool dry place

## Ronifur

Antibiotic tablets against trichomoniasis, hexamitiasis, infections of the gastrointestinal tract.

**Composition for one tablet 150mg:** Ronidazole: 7.5mg, Furaltadone HCl: 7.5mg.

**Target species:** Pigeons and decorative birds.

**Indications:** For treatment and prophylaxis of trichomoniasis, gangrene, E-coli, paratyphoid, coccidiosis, the gastro-intestinal infections, diarrhea.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Pigeons and decorative birds: 1 - 2 tablets once a day for 5-10 days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Amoxi Tab

Antibiotic tablets against bacterial and polymicrobial diseases

**Composition for one tablet 280mg:** Amoxicillin trihydrate: 10mg, Complex of vitamins

**Target species:** Pigeons.

**Indications:** For prophylaxis and treatment of bacterial or respiratory infections, enterotoxic diseases, E-coli, chronic respiratory infections, polymicrobial diseases with unknown etiology.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** Small pigeons: ½ tablet per day for 5-7 consecutive days

Grown-up pigeons: 1 tablet per day for 5-7 consecutive days

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Tetra-amoxy-neo Tab

Antibiotic complex for treatment and prophylaxis of bacterial and chronic respiratory infections. Tablets.

**Composition for one tablet 280mg:** Oxytetracycline hydrochloride: 10mg, Amoxicillin trihydrate: 9mg, Neomycin sulphate: 9mg, Complex of vitamins.

**Target species:** Pigeons.

**Indications:** For treatment and prophylaxis of bacterial or chronic respiratory infections, enterotoxic diseases, E-coli, polymicrobial diseases with unknown etiology.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** 1 - 2 tablets per kg body weight once a day for 5-7 days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Tricho Cocci Vit

Antibiotic tablets and vitamin complex for treatment of coccidiosis, trichomonosis and hexamitose.

**Composition for one tablet 150mg:** Ronidazole: 10mg, Amprolium HCL: 10mg, Vitamin A, Vitamin D3, Vitamin E, Vitamin K, Vitamin C, Vitamin B1, Vitamin B2, Vitamin B6, Vitamin B12.

**Target species:** Pigeons.

**Indications:** Tricho-Cocci Vit Tab. is active against coccidiosis, trichomonosis and hexamitose. The vitamin complex helps to keep pigeons in good shape.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally.

**Dosage:** 1 tablet once a day for 2-3 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 50pcs 100pcs

Store below 28°C in cool dry place

## Rodifen Combi Vit

Antibiotic and vitamin complex against internal parasites. Tablets.

**Composition for one tablet 200mg:** Ronidazole: 18mg, Diclazuril: 2.5mg, Fenbendazole: 10mg

**Target species:** Pigeons.

**Indications:** Active against coccidiosis, trichomonosis and other internal parasites. The vitamin complex helps to keep pigeons in good shape.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Orally, on an empty stomach.

**Dosage:** 1 tablet once a day on an empty stomach! Repeat treatment after 3-4 weeks. Treat all birds on the premises! Disinfect the premises!

If there is a risk of infection treat birds at the beginning of the reproductive period, with the incubation of new birds.

**Withdrawal period:** Not applicable.





Package size: **40g**

Store below 28°C in cool dry place

## Neobacin

Antibiotic for treatment of skin infections after surgical procedures, cuts, burns, lacerations, wet wounds. Cutaneous powder.

**Composition in 1g:** Neomycin sulphate : 5 000 UI, Zinc bacitracin: 250 UI

**Excipients:** Kaolin to 1g

**Target species:** Pigeons, decorative birds, decorative rabbits, guinea pigs.

**Indications:** An antibiotic combination for external use. The two highly effective antibiotics have mutually reinforcing effects. The product is not being inactivated by blood and secretions and is suitable for wounds. Used for treatment of skin infections after surgical procedures, also for cuts, burns, lacerations, wet wounds.

**Route of administration:** Externally on skin.

**Dosage:** Apply thin layer over the infected area 2 to 4 times a day.

**Withdrawal period:** Not applicable.



Package size: **40g**

Store below 28°C in cool dry place

## Sulphathiazole

Antibiotic for treatment of open wounds, burns, decubitus, ulcers, etc. Cutaneous powder.

**Composition in 100g:** Sulphathiazole

**Excipients:** None

**Target species:** Pigeons and decorative birds.

**Indications:** Antibiotic with an antibacterial and bacteriostatic effect against streptococci, staphylococci. Boosts the synthesis of folic acid, which is necessary for the synthesis of purine nucleotides in the protein cell. Side reactions are rarely observed.

**Route of administration:** Externally on skin.

**Dosage:** Apply thin layer over the infected area 2 to 4 times a day.

**Withdrawal period:** Not applicable.



## Oxytetra Ointment

Antibiotic against bacterial skin infections. Cutaneous ointment.

**Composition in 100g:** Oxytetracycline HCl: 1g

**Target species:** Pigeons and decorative animals (hamsters, guinea pigs, decorative rabbits).

**Indications:** For local treatment of skin infections caused by oxytetracycline-sensitive microorganisms, surgical wounds, ulcers.

**Route of administration:** Externally on skin.

**Dosage:** Clean the infected area and apply 2 times a day.

**Withdrawal period:** Not applicable.

Package size: **50g**

Store below 28°C in cool dry place



## Sulphathiazole

For treatment of wounds and skin infections. Cutaneous ointment.

**Composition in 100g:** Norsulfazolum: 5g, Oleum Yekoris: 5g.

**Excipient:** Dextrose monohydrate to 100g.

**Target species:** Pigeons.

**Indications:** For treatment of infected wounds, operative wounds, abrasions, skin infections, etc. Has an antimicrobial effect on streptococci, staphylococci, pneumococci, coliform bacteria, additionally it has an anti-allergic and antiphlogistic effect.

**Route of administration:** Externally on skin.

**Dosage:** Clean the infected area and apply by a spatula in the morning and evening for 4-5 days.

*The product should not be administered to poultry intended for human consumption.*

**Withdrawal period:** Not applicable.

Package size: **50g 500g 1Kg**

Store below 28°C in cool dry place



## Trio Ointment

**For treatment of wounds, ulcers, eczema, dermatitis. Cutaneous ointment.**

**Composition in 100g:** Gentamicin sulfate: 1%, Neem oil: 1%, Clove oil: 1%, Vitamin A: 50 000 000UI

**Target species:** Pigeons and decorative animals.

**Indications:** Gentamicin has antimicrobial and anti-inflammatory properties. Neem oil has an antiparasitic effect, and Clove oil has an antifungal and analgesic effect. Vitamin A helps the epithelialization of the skin. It is used for freshly infected wounds, ulcers, eczema, dermatitis, itching of unknown origin and has antiparasitic and antifungal effect.

**Route of administration:** Externally on skin.

**Dosage:** Apply twice - in the morning and in the evening.

**Withdrawal period:** Not applicable.

**Package size:** 50g

Store below 28°C in cool dry place



## Neobacin

**Antibiotic emulsion against skin diseases**

**Composition in 100ml:** Neomycin sulphate: 5 000 UI, Zinc bacitracin: 250 UI, Peppermint oil and clove oil

**Excipient:** to 100 ml

**Target species:** Pigeons and decorative animals (guinea pigs, decorative rabbits).

**Indications:** The combination of the two antibiotics, as well as the complementary action of the two removes itching and pain and has an antifungal effect. Suitable for insect bites, pain and itching in the ears. The preparation enhances the recovery process. Used for treatment of skin damage, cuts, lacerations, postoperative treatments.

**Route of administration:** Externally on skin.

**Dosage:** Shake before use. Clean the infected area and apply 2-3 times a day until complete recovery.

**Withdrawal period:** Not applicable.

**Package size:** 100ml

Store below 28°C in cool dry place



## Neocubatul Spray

**Cutaneous spray with insecticidal, antifungal and antiseptic effect**

**Composition in 100ml:** Pix Liquida: 15%

**Target species:** Pigeons and exotic animals and pets - hamsters, mini rabbits, guinea pigs, ornamental birds and others.

**Indications:** The spray has insecticidal, antifungal and antiseptic effect. It forms protective coating against irritation and enhances the regenerative processes. Used for wounds, burns, castrations, dermatitis, wet and dry eczema, dermatomycosis, for treatment of umbilical cord and has a repellent action.

**Route of administration:** Externally on skin.

**Dosage:** Spray affected area from a distance of 15-20 cm, for 2-3 consecutive days.

**Withdrawal period:** Not applicable.

**Package size:** 100ml 140ml

Store below 28°C in cool dry place



## Oxytetra Spray

**Solution with antimicrobial, anti-allergic and anti-inflammatory effect.**

**Composition in 100ml:** Oxytetracycline hydrochloride: 2.00g, Hydrocortisone acetate: 0.10g

**Excipient:** Glycerol, Citric acid, Ethyl alcohol 70%

**Target species:** Pigeons and decorative animals (guinea pigs, decorative rabbits).

**Indications:** Aerosol with antimicrobial, anti-allergic and anti-inflammatory effect. It is highly effective in the treatment of infected skin, mucosal and traumatic damages, random and operating wounds and dermatitis.

**Route of administration:** Externally on skin.

**Dosage:** Clean the affected area thoroughly of necrotic tissues, exudates and other contaminants, then spray for 1-2 seconds from a distance of 15-20 cm. Re-treat every 12 hours until complete recovery of the affected area.

**Withdrawal period:** Not applicable.

**Package size:** 200ml

Store below 28°C in cool dry place



## Neobacin Spray

Against skin infections caused by traumatic injuries, cuts, burns, lacerations, wet wounds, surgical interventions.

**Composition in 100ml:** Neomycin sulphate: 5 000 UI, Zinc bacitracin: 250 UI

**Target species:** Pigeons and decorative animals (guinea pigs, decorative bunnies).

**Indications:** The two highly effective antibiotics have mutually reinforcing effects. The product is not inactivated by blood and secretions and is very suitable for wounds. Used to treat skin infections after traumatic injuries, cuts, burns, lacerations, wet wounds. It is also used after surgical interventions.

**Route of administration:** Externally on skin.

**Dosage:** Clean the affected area thoroughly of necrotic and foreign tissues, exudates and other contaminants, then spray for 1-2 seconds from a distance of 15-20 cm, 2-3 times a day until complete recovery.

**Withdrawal period:** Not applicable.

**Package size:** 100ml

Store below 28°C in cool dry place



## Eczeform C

Cutaneous spray solution against skin diseases, eczema, mycosis, septic wounds, ulcers, abscesses, decubitus and others.

**Composition in 100ml:** Sulfathiazolum: 2.5g, Acidum salicylicum: 1.6g, Resorcinum: 1.6g, Oleum ricini: 6.3g

**Excipient:** Glycerinum, Ethyl alcohol 70% - to 100ml.

**Target species:** Pigeons.

**Indications:** For treatment of skin diseases, eczema, mycosis, septic wounds, ulcers, abscesses, decubitus and others.

To be used only for target animals.

**Route of administration:** Cutaneously, locally, by spraying.

**Dosage:** The skin of the diseased area is cleansed in advance and rubbed with a swab twice a day.

**Withdrawal period:** Not applicable.

**Package size:** 50ml 100ml

Store below 28°C in cool dry place



## Oti Farm

Ear drops against inflammations, fungal and bacterial infections.

**Composition in 1ml:** Oxytetracycline HCl: 20mg/ml, Prednisolone acetate: 5.0mg/ml, Ichthammol: 100mg/ml

**Target species:** Pigeons, hamsters, guinea pigs, decorative bunnies and other exotic animals.

**Indications:** Against otitis externa caused by fungal or bacterial infections with antifungal, bactericidal and anti-inflammatory action. Reduces itching and acts as an analgesic.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Auricular use.

**Dosage:** 3-5 drops in the affected ear, twice a day with subsequent massage.

**Withdrawal period:** Not applicable.

**Package size:** 10ml

Store below 28°C in cool dry place



## Oti Lucil

Ear drops with antimicrobial, anti-allergic and anti-inflammatory effect.

**Composition in 100ml:** Oxytetracycline HCl: 0.15g, Cypermethrine: 0.20g, Salicylic acid: 0.22g

**Excipient:** Glycerin, Carnation oil extract, Mint oil.

**Target species:** Pigeons, hamsters, guinea pigs, decorative bunnies and other exotic animals.

**Indications:** Against inflammations, ear scabies, itching - caused by fungal inflammation, parasites, as well as itching of unknown origin. Against otitis externa caused by fungal or bacterial infections. Reduces itching and acts as an analgesic.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Auricular use.

**Dosage:** 3-5 drops in the infected ear, twice a day with subsequent massage.

**Withdrawal period:** Not applicable.

**Package size:** 10ml

Store below 28°C in cool dry place





## Iodine - Potassium - Iodine Ointment

Cutaneous ointment with antiseptic, antimicrobial, antimycotic, acaricidal action.

**Composition in 100g:** Pure iodine: 5.0g , Potassium iodide: 10.0g

**Target species:** Pigeons.

**Indications:** : It has an antiseptic, antimicrobial, antimycotic, acaricidal action. The ointment causes hyperthermia that helps to spread infiltrates and promotes granulation. Iodine penetrates deeply and can have a resorptive effect. It is used in inflammatory skin changes in wounds and dermatomycosis. The product should not be administered to poultry intended for human consumption.

**Route of administration:** : Externally on skin, locally.

**Dosage:** Rub into the skin 1-2 times a day.

**Withdrawal period:** Not applicable.

**Package size:** 50g

Store below 28°C in cool dry place



## Propolis

Natural product with antimicrobial, antiviral and anti-radiation effect. Cutaneous ointment.

**Composition in 100g:** Propolis 10g

**Target species:** Pigeons and decorative animals (guinea pigs, decorative rabbits.)

**Indications:** Propolis is a mixture of resins, wax and pollen. It contains vitamins, essential oils, mineral salts, trace elements, enzymes, hormones and ferments. Propolis has antimicrobial, antiviral, anti-mold, anti-radiation effect, strong antibiotic and antimicrobial action. Acts as a disinfectant and analgesic. Used in skin changes with unclear origin - dermatitis, eczema, hair loss, wounds and ulcers.

**Route of administration:** Externally on skin.

**Dosage:** Apply to the infected area 1-2 times a day.

**Withdrawal period:** Not applicable.

**Package size:** 50g 500g 1Kg

Store below 28°C in cool dry place



## Eczefarm A

Antiseptic for wet and dry eczema. Cutaneous ointment.

**Composition in 100g:** Salicylic acid: 1.3g, Purified sulfur: 6.8g

Coniferous tar: 23g

**Excipients:** Paraffin white soft to 100g

**Target species:** Pigeons.

**Indications:** Used for wet and dry eczema and in inflammatory skin changes, has antiseptic, antifungal, insecticidal, acaricidal, keratoplastic properties. It causes hyperemia of the skin and creates an opportunity to limit the inflammation.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Externally on skin.

**Dosage:** Apply 1-2 times a day for 5-6 days. consecutive days.

**Withdrawal period:** Not applicable.

**Package size:** 50g

Store below 28°C in cool dry place



## Eczefarm B

Antiseptic for wet eczema and dermatitis. Cutaneous ointment.

**Composition in 100g:** Zinc oxide: 14g, Bismuth: 14g, Ichthyol: 1.4g .

**Excipient:** White Soft Paraffin to 100g.

**Target species:** Pigeons.

**Indications:** Used for treatment of wet eczema and dermatitis. Reduces pain and itching, has antiseptic, antifungal, insecticidal, acaricidal, keratoplastic, astringent and antiphlogistic properties. Accelerates wound healing, has drying and absorbing effect, softens the skin.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Externally on skin.

**Dosage:** Apply locally 1-2 times a day for 5-6 days.

**Withdrawal period:** Not applicable.

**Package size:** 50g

Store below 28°C in cool dry place



## Granufarm Spray

For treatment of newly infected wounds, decubital wounds and ulcers, eczema, dermatitis. Cutaneous spray.

**Composition in 100ml:** Ethacridini lactat: 0.37g, Flavacridini hydrochloridum: 0.37g, Methylen blau: 0.74g, Acidum salicylicum: 2.22g

**Excipient:** Ethyl alcohol 70% to 100 ml.

**Target species:** Pigeons and decorative animals.

**Indications:** For treatment of newly infected wounds, decubital wounds and ulcers, eczema, dermatitis.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Externally on skin.

**Dosage:** Clean the infected area and apply locally twice a day.

**Withdrawal period:** Not applicable.

**Package size:** 50ml 100ml

Store below 28°C in cool dry place



## Iodine Tincture 5 % Spray

Disinfection solution. Cutaneous spray.

**Composition in 100ml:** Iodine: 5.0g, Potassium iodide: 2.0g

**Excipient:** Purified water and Ethanol 95%.

**Target species:** Pigeons.

**Indications:** The iodine tincture has a bactericidal, sporicidal, virucidal, antimycotic, acaricidal (in itch-mites, *Sarcoptes scabiei*), anthelmintic and antiprotozoal effect. It causes hyperemia at the site of administration. For the disinfection of the operative field, for the treatment of wounds (around them).

**Route of administration:** Externally on skin.

**Dosage:** Clean the infected area and apply locally twice a day

**Withdrawal period:** Not applicable.

**Package size:** 50ml 100ml

Store below 28°C in cool dry place



**Package size:**

20ml 50ml 100ml 200ml 1L 5L

Store below 28°C in cool dry place

## Iodine Tincture 5%

Disinfection solution. Cutaneous solution.

**Composition in 1ml:** Iodine: 5.0g, Potassium iodide: 2.0g.

**Excipients:** Purified water and Ethanol 95%.

**Target species:** Pigeons.

**Indications:** The iodine tincture has a bactericidal, sporicidal, virucidal, antimycotic, acaricidal (in itch-mites, *Sarcoptes scabiei*), anthelmintic and antiprotozoal effect. For disinfection of the operative field, for treatment of wound surrounding area.

**Route of administration:** Externally on skin.

**Dosage:** Clean the infected area and apply locally twice a day.

**Withdrawal period:** Not applicable.



**Package size:**

50ml 100ml 200ml 1L

Store below 28°C in cool dry place

## Granufarm

For treatment of freshly infected wounds, decubital wounds and ulcers, eczema, dermatitis. Cutaneous solution.

**Composition in 100ml:** Ethacridini lactat: 0.37g, Flavacridini hydrochloridum: 0.37g, Methylenblau: 0.74g, Acidum salicylicum: 2.22g

**Excipient:** Ethyl alcohol 70% to 100 ml.

**Target species:** Pigeons and decorative animals.

**Indications:** For treatment of freshly infected wounds, decubital wounds and ulcers, eczema, dermatitis.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Auricular use.

**Dosage:** Clean the infected area and apply locally twice a day.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Calcivit

Calcium and phosphorus supplement. Tablets.

**Composition for one tablet 750mg:** Calcium: 270mg, Phosphorus: 170mg, Lactose, Starch, Vitamin A: 80 UI, Vitamin D3: 8 UI, Vitamin C: 0.40mg, Nicotinamide: 0.40mg, Niacin: 0.208mg, Vitamin B2 (Riboflavine): 0.040mg, Calcium-D-pantothenate: 0.040mg, Vitamin B6: 0.016mg, Vitamin K3: 0.016mg, Vitamin E: 0.0112mg, Vitamin B1: 0.0104mg, Folic acid: 0.0016mg, Biotin: 0.00056mg, Vitamin B12: 0.000040mg;

**Target species:** Pigeons and decorative birds.

**Indications:** For calcium and phosphorus deficiencies.

**Route of administration:** Orally.

**Dosage:** 1 tablet daily.

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 100pcs

Store below 28°C in cool dry place

## Polyvit

For prophylaxis of vitamin deficiency. Tablets

**Composition for one tablet 750mg:** Vitamin A: 80 UI, Vitamin D3: 8 UI, Vitamin C: 0.40mg, Nicotinamide: 0.40mg, Niacin: 0.208mg, Vitamin B2 (Riboflavine): 0.040mg, Calcium-D-pantothenate: 0.040mg, Vitamin B6: 0.016mg, Vitamin K3: 0.016mg, Vitamin E: 0.0112mg, Vitamin B1: 0.0104mg, Folic acid: 0.0016mg, Biotin: 0.00056mg, Vitamin B12: 0.000040mg;

**Target species:** Pigeons and decorative birds.

**Indications:** Vitamin complex with multiple effect that increases the immunobiological activity, affects moulting, reproduction, coupling, fertilization and hatching.

**Route of administration:** Orally.

**Dosage:** 1/4 tablet daily.

**Withdrawal period:** Not applicable.



Package size:

10ml 20ml 50ml 100ml 1L 5L

Store below 28°C in cool dry place

## Maxi Vit - Vitamin supplement

Oral solution.

**Composition in 100ml:** Vitamin A (retinol acetate): 20 000 000 UI, Vitamin D3 (cholecalciferol): 500 000 IU, Vitamin E (alpha tocopherol acetate): 8000 IU, Vitamin B1 (Thiamini hydrochloride): 1250mg, Vitamin B2 (Riboflavine): 1250mg, Vitamin B6 (Pyridoxine hydrochloride): 500mg, Vitamin B12 (Cyanocobalamin): 5mg, Vitamin K3 (Menadione natrii Bisulfite): 500mg, Vitamin H: 15mg, Vitamin PP: 10 000mg, Choline chloride: 100 000mg

**Target species:** Pigeons and decorative birds.

**Indications:** Provides necessary vitamins to accelerate growth, increase productivity and promote efficient metabolic processes.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1 ml in 3-4 liters of water for 10-15 days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Vitatonic

Powder for oral solution. Vitamin complex.

**Composition in 100g:** Vitamin A: 100 000 IU, Vitamin D3: 10 000 IU, Vitamin C: 500mg, Nicotinamide: 500mg, Niacin: 260mg, Vitamin B2: 50mg, Calcium-D-pantothenate: 50mg, Vitamin B6: 20mg, Vitamin K3: 20mg, Vitamin E: 14mg, Vitamin B1: 13mg, Folic acid: 2mg, Biotin: 0.7mg, Vitamin B12: 0.05mg, Dextrose to 100g.

**Target species:** Pigeons and decorative birds.

**Indications:** Vitamin complex with multiple beneficial effect on the body. Affects the absorption of nutrients, and thus helps the body to convert food into energy. Increases the immunobiological reactivity and resistance of the body to adverse environmental factors, as well as its resistance to different diseases. Acts favorably in stress conditions, as well as in conditions associated with increased vitamin needs - pregnancy, lactation, egg production, after treatment of infectious diseases, etc.

**Route of administration:** Internally, dissolved in the drinking water or with the feed.

**Dosage:** 1-2g / l of water for 3-5 consecutive days or 2.5 g / 10 kg feed for 3-5 consecutive days.

**Withdrawal period:** Not applicable.





Package size:

10ml 20ml 100ml 500ml 1L 5L

Store below 28°C in cool dry place

## Maxi Vit - Vitamin complex

**Solution for oral administration. Accelerates growth, increases the productivity and stability of the animal, supports the metabolic processes.**

**Composition in 100ml:** Vitamin A: 20 000 000 IU, Vitamin D3: 500 000 IU, Vitamin E: 8000 IU, Vitamin B1: 1250 mg, Vitamin B2: 1250 mg, Vitamin B6: 500 mg, Vitamin B12: 5 mg, Vitamin K3: 500 mg, Vitamin H: 15 mg, Vitamin PP: 10 000 mg, Choline chloride: 100 000 mg

**Target species:** Birds, swine, calves, pigs, lambs.

**Indications:** The vitamin complex provides all necessary vitamins for birds, swine, calves, lambs, etc. in cases of vitamin deficiency and parasite caused diseases.

**Route of administration:** Orally with drinking water.

**Dosage:** Dissolved in drinking water for 10-15 days. For birds-1ml. in 3, 4 l. of water; For pigs-0,5 to 1ml. per animal, for calves - 2 ml per animal.

**Withdrawal period:** Not applicable.



## Vitamin E hyrdozole

**Liquid preparation. Emulsion for oral solution with anti - sterile action.**

**Composition in 100 ml:** Vitamin E (alpha tocopheryl acetate): 6.000g

**Excipients:** Preservative E 211, Emulsifier E 433

**Target species:** Birds, calves, pigs, lambs, dogs and rabbits.

**Indications:** Vitamin E has anti- sterile action in both female and male animals and birds. It supports the metabolic process and facilitates the functions of the glands, nerves and muscles, strengthens the immune system. It is used for prophylaxis of vitamin E deficiency.

**Route of administration:** Internally, with the drinking water.

**Dosage:** For birds-1ml / 1L water; for calves, pigs, lambs, rabbits, dogs- 0,1 ml/ kg.

**Withdrawal period:** Not applicable.

Package size: 50ml 100ml 1L

Store below 28°C in cool dry place



## AD3E

**Emulsion for fertility and reproduction support.**

**Composition in 1L:** Vitamin A (E672) 50 000 000 IU, Vitamin D3 (E671): 5 000 000 IU, Vitamin E: 700, All-rac-alpha-tocopheryl acetate 30 000 mg

**Excipients:** Glycerol polyethylene glycol ricinoelate (E484): 250 000 mg Sorbitol (E420): 150 000 mg, Moisture: 50.99%

**Target species:** Birds, pigs, rabbits, horses, cattle.

**Indications:** For reproduction and fertility support.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** Birds: 0,2-10 ml/ 1L. water; calves: 10ml./ 1L. water or 8ml. directly through mouth / day; milk cows/ cattle/ horses: 10 ml./ 1L water, or 20 ml. directly through mouth / day; pigs- 20 ml/ 1L. water, or 1-8 ml. directly through mouth/ day; rabbits- 20 ml/ 1L water, or 2ml directly / day

**Withdrawal period:** Not applicable.

Package size: 50ml 100ml 500ml 1L

Store below 28°C in cool dry place



Package size: **40g**

Store below 28°C in cool dry place

## Fipronil

Against poultry red mites, fleas, flies, ants, cockroaches, and other ectoparasites. Cutaneous powder.

**Composition in 100g:** Fipronil: 1.0g.

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** Against poultry red mites, fleas, flies, ants, cockroaches, and other infections transmitting ectoparasites.

**Route of administration:** Sprinkle deeply against the hair until it reaches the skin. Treat the head, ears, neck, back, limbs and paws. Also powder the premises and sheds.

**Dosage:** Apply thin layer over the infected area 2 to 4 times a day.

**Withdrawal period:** Not applicable.



Package size: **40g**

Store below 28°C in cool dry place

## Trio Kill Forte

Against ticks, lice, fleas, biting lice and their eggs. Cutaneous powder.

**Composition in 100g:** Cypermethrin: 0.5g, Nuvan: 0.5g, Fipronil: 0.1g

**Excipients:** None

**Target species:** Pigeons, guinea pigs and decorative rabbits.

**Indications:** Ectoparasite powder against ticks, lice, fleas, biting lice and their eggs.

**Route of administration:** Externally on skin.

**Dosage:** Ready to use powder. Treat the head, ears, neck, back, limbs and paws of the animal.

**Withdrawal period:** Not applicable.



## Trio Kill Forte

Against red mites, fleas, flies, ants and other ectoparasites. Cutaneous spray.

**Composition in 100g:** Cypermethrin: 0.2g, Nuvan: 0.1g, Fipronil: 0.01g

**Target species:** Pigeons, exotic birds and decorative animals. The product should not be administered to poultry intended for human consumption.

**Indications:** Against poultry red mites, fleas, flies, ants, lice and other infection transmitting ectoparasites.

The product should not be administered to poultry intended for human consumption.

**Route of administration:** Externally on skin.

**Dosage:** Spray at a distance of 15 cm under the wings and on the back of the pets.

**Withdrawal period:** Not applicable.

Package size: **200ml**

Store below 28°C in cool dry place



## Deltamethrin Colombo

For bathing against red mites, ticks, lice, fleas and all infection transmitting ectoparasites. Cutaneous spray.

**Composition in 100g:** Deltamethrin: 0.1g.

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** Against poultry red mites, ticks, lice, fleas and all infection transmitting ectoparasites.

**Route of administration:** Externally on skin.

**Dosage:** Spray at a distance of 15 cm under the wings and on the back of the pets.

**Withdrawal period:** Not applicable.

Package size: **200ml** **500ml**

Store below 28°C in cool dry place



Package size:

10ml 20ml 50ml 100ml 1L

Store below 28°C in cool dry place

## Fantomas Forte

Against poultry red mites, fleas, flies, ants, ticks, lice and other ectoparasites. Concentrate for cutaneous solution.

**Composition in 100g:** Deltamethrin: 2.5g, Cypermethrin: 8g, Piperonyl butoxide: 5g

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** For bathing against poultry red mites, fleas, flies, ants and other infections transmitting ectoparasites.

**Route of administration:** Externally, dissolved in bathing/spraying water. Spray all the premises with the same solution.

**Dosage:** 1 ml/1 L water.

**Contraindications:** Atropine

**Withdrawal period:** Not applicable.



Package size:

10ml 20ml 50ml 100ml 1L

Store below 28°C in cool dry place

## Fantomas

Against poultry red mites, fleas, flies, ants, ticks, lice and other ectoparasites. Concentrate for cutaneous solution.

**Composition in 100g:** Deltamethrin: 2.5g, Cypermethrin: 5.0g, Piperonyl butoxide: 2.5g

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** For bathing against poultry red mites, fleas, flies, ants and other infections transmitting ectoparasites.

**Route of administration:** Externally, dissolved in bathing/spraying water. Spray all the premises with the same solution

**Dosage:** 1 ml/1 L water.

**Contraindications:** Atropine

**Withdrawal period:** Not applicable.

## Fipronil Spot On

Against fleas, lice, ticks, poultry red mites, biting lice and other ectoparasites. Drops for cutaneous use.

**Composition in 100g:** Fipronil: 2g

**Target species:** Pigeons.

**Indications:** Insecticidal product against fleas, lice, ticks, poultry red mites, biting lice and other ectoparasites.

**Route of administration:** Externally on skin.

**Dosage:** Ready to use product.

**Withdrawal period:** Not applicable.



Package size: 10ml

Store below 28°C in cool dry place

## Fipronil Spray

Solution against red mites, ticks, lice, fleas and all ectoparasites. Cutaneous spray

**Composition in 1g:** Deltamethrin: 2.5mg

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** Against poultry red mites, ticks, lice, fleas and all infection transmitting ectoparasites.

**Route of administration:** Externally on skin.

**Dosage:** Spray at a distance of 15 cm under the wings and at the back of the pets. Spray the premises with the same solution.

**Withdrawal period:** Not applicable.



Package size: 100ml

Store below 28°C in cool dry place





## Insectostop Colombo

**Against poultry red mites, fleas, flies, ants, ticks, lice and other ectoparasites. Concentrate for cutaneous solution**

**Composition in 100g:** Dichlorvos: 35 g

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** For bathing against poultry red mites, fleas, flies, ants and other infections transmitting ectoparasites.

**Route of administration:** Externally, dissolved in bathing/spraying water. Spray the premises with the same solution.

**Dosage:** 1 ml/1 L water.

**Contraindications:** Atropine

**Withdrawal period:** Not applicable.

Package size:

9ml	10ml	20ml	50ml	100ml
500ml	1L	5L		

Store below 28°C in cool dry place



## Nuvan Spray

**Against poultry red mites, fleas, flies, ants, ticks, lice and other infection transmitting ectoparasites. Cutaneous spray.**

**Composition in 100g:** Dichlorvos: 0.2g

**Target species:** Pigeons, decorative birds and decorative animals.

**Indications:** Against poultry red mites, fleas, flies, ants and other infection transmitting ectoparasites.

*The product should not be administered to poultry intended for human consumption.*

**Route of administration:** Externally on skin. Spray the premises with the same solution.

**Dosage:** Ready to use solution.

**Withdrawal period:** Not applicable.

Package size: 200ml 500ml

Store below 28°C in cool dry place



## Nuvan Colombo

**For bathing against external parasites. Concentrate for cutaneous solution.**

**Composition in 100g:** Dichlorvos: 35g.

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** For bathing against poultry red mites, fleas, flies, ants and other infections transmitting ectoparasites - ticks, lice and other.

**Route of administration:** Externally, dissolved in bathing/spraying water. Spray the premises with the same solution.

**Dosage:** 1 ml/1 L water.

**Withdrawal period:** Not applicable.

Package size:

10ml	20ml	100ml	500ml	1L
------	------	-------	-------	----

Store below 28°C in cool dry place



## Tetrazin Forte

**Triple protection against poultry red mites, fleas, flies, and other ectoparasites. Concentrate for cutaneous solution.**

**Composition in 100g:** Cypermethrin: 10g, Nuvan: 3.0g, Fipronil: 0.1g.

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** For bathing and spraying against poultry red mites, fleas, flies, and other infection transmitting ectoparasites.

**Route of administration:** Externally, dissolved in bathing/spraying water.

**Dosage:** 1 ml/1 L water.

**Withdrawal period:** Not applicable.

Package size:

10ml	20ml	100ml	500ml	1L
------	------	-------	-------	----

Store below 28°C in cool dry place



Package size:

10ml 20ml 100ml 500ml 1L 5L

Store below 28°C in cool dry place

## Trio Kill Forte

Against red mites, fleas, flies, ants and other ectoparasites.  
Concentrate for cutaneous solution.

**Composition in 100g:** Cypermethrin: 3.5g, Tetramethrin: 1.5g, Piperonyl butoxide: 4.7g

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** For bathing against external parasites - poultry red mites, fleas, flies, ants and other infections transmitting ectoparasites.

**Route of administration:** Externally, dissolved in bathing/spraying water.

**Dosage:** 1 ml/1 L water.

**Contraindications:** Atropine

**Withdrawal period:** Not applicable.



## Ectolucyl Spray

Cutaneous spray with antiparasitic and antibacterial effect.

**Composition in 100g:** Oxytetracycline hydrochloride: 2.00g, Cypermethrin: 0.20g, Castor oil

**Target species:** Pigeons and decorative animals (guinea pigs, decorative rabbits).

**Indications:** The product combines the antibacterial action of oxytetracycline with the antiparasitic effect of cypermethrin, softening and epithelizing action of castor oil. Oxytetracycline hydrochloride is a broad-spectrum antibiotic. It acts on Gram-positive and Gram-negative microorganisms. Ectolucil spray is used in wounds and ear scabies for decorative bunnies, hairless spots, caused by *Sarcoptes scabiei*.

**Route of administration:** Externally on skin.

**Dosage:** Spray the affected areas at a distance of 10-15 cm once a day for 2-3 consecutive days. Repeat after 15 days.

**Withdrawal period:** Not applicable.

Package size:

100ml

Store below 28°C in cool dry place



Package size:

10ml 20ml 100ml 500ml 1L 5L

Store below 28°C in cool dry place

## Deltamethrin Colombo

For bathing against external parasites. Concentrate for cutaneous solution.

**Composition in 100g:** Deltamethrin: 5g.

**Target species:** Pigeons, exotic birds and decorative animals.

**Indications:** For bathing against external parasites. With the same solution spray the rooms, sheds and all places where the ectoparasites could be found.

**Route of administration:** Externally, dissolved in bathing/spraying water.

**Dosage:** 1 ml / 1 L water.

**Withdrawal period:** Not applicable.



Package size:

10ml 20ml 100ml 500ml 1L

Store below 28°C in cool dry place

## Carbaryl Colombo

Cutaneous solution with 10 % Carbaryl against red mites, ticks, lice, flies, ants, cockroaches and other ectoparasites.

**Composition in 100g:** 10% Carbaryl

**Target species:** Pigeons, exotic birds, exotic animals.

**Indications:** Against red mites, ticks, lice, flies, ants, cockroaches and other ectoparasites.

**Route of administration:** Externally, for bathing against ectoparasites. With the same solution spray under the wings and other body parts where ectoparasites can be found.

**Dosage:** 1 ml/1 L water.

**Withdrawal period:** Not applicable.



Package size:

10ml 50ml 100ml 1L

Store below 28°C in cool dry place

## Protector Forte

Cutaneous solution against ectoparasites.

**Composition in 100g:** Deltametrin: 2.5%, Fipronil: 0.1%, Piperoniol butoxide: 2%

**Target species:** Pigeons, exotic animals and birds.

**Indications:** For bathing against infection transmitting ectoparasites- red mites, fleas, flies, ticks, ants, cockroaches. With the same solution spray the premises where the animals live.

**Route of administration:** Externally on skin, or spraying the premises.

**Dosage:** 1ml/ 2 l water.

**Contraindications:** Atropine

**Withdrawal period:** Not applicable.



Package size:

10ml 50ml 100ml 1L

Store below 28°C in cool dry place

## Aytcol Colombo

Cutaneous solution against ectoparasites.

**Composition in 100g:** Deltametrin: 5%, Dichlorvos: 0.5%

**Target species:** Pigeons, exotic animals and birds.

**Indications:** For bathing against infection transmitting ectoparasites- red mites, fleas, flies, ticks, ants, cockroaches. With the same solution spray the premises where the animals live.

**Route of administration:** Externally on skin, or spraying the premises.

**Dosage:** 1ml/ 1L water

**Withdrawal period:** Not applicable.



Package size:

10ml 20ml 100ml 1L

Store below 28°C in cool dry place

## Aytcol Forte Colombo

Cutaneous solution against ectoparasites

**Composition in 100g:** Deltametrin: 5%, Dichlorvos 10%

**Target species:** Pigeons, exotic animals and birds.

**Indications:** For bathing against infection transmitting ectoparasites- red mites, fleas, flies, ticks, ants, cockroaches. With the same solution spray the premises where the animals live.

**Route of administration:** Externally on skin, or spraying the premises.

**Dosage:** 1ml/ 2L water.

**Contraindications:** Atropine

**Withdrawal period:** Not applicable.



Package size:

20pcs 30pcs 50pcs 100pcs

Store below 28°C in cool dry place

## Levafarm

For treatment of gastrointestinal and lung nematodes.

**Composition for one tablet 150mg:** Levamisole hydrochloride: 23mg.

**Excipients:** Lactose, Starch, Magnesium stearate, Talc.

**Target species:** Pigeons and decorative rabbits.

**Indications:** : For treatment of gastrointestinal and lung nematodes in pigeons and decorative rabbits.

**Route of administration:** Orally.

**Dosage:** Twice a year: ½ - 1 tablet. Do not feed the pigeons 24 hours before and 2-3 hours after the treatment. In case of severe infestations, repeat after 3 weeks. Disinfect the pigeon-house! The product should not be used in poultry intended for human consumption and in birds that feed their little ones.

**Withdrawal period:** Not applicable.





Package size:

20ml 100ml 200ml 1L 5L

Store below 28°C in cool dry place

## Piperazine 20%

For treatment of ascariasis and capillariasis. Oral solution.

**Composition in 100ml:** Piperazine hexahydrate 20g.

**Excipients:** Sulphuric acid, Methyl parabene and Aqua purificata.

**Target species:** Pigeons.

**Indications:** For treatment of ascariasis and capillariasis.

**Route of administration:** Internally with soft food. Feed birds in evenings only! Do not feed birds 12-15 hours before treatment!

**Dosage:** 2.0-3.0 ml / kg weight.

12 to 14 hours after treatment excrements should be mechanically cleaned, floors should be disinfected.

*The treatment should be repeated in 2 months.*

**Contraindications:** Atropine

**Withdrawal period:** Not applicable.



Package size:

10ml 20ml 100ml 500ml 1L 5L

Store below 28°C in cool dry place

## Pharmacox

For treatment of coccidia, toxoplasma, sarcocystis, taylorii.

**Composition in 100ml:** Toltrazuril: 2.5 g

**Target species:** Pigeons and decorative birds.

**Indications:** Toltrazuril is a synthetic compound from the group of symmetrical triazinones. It acts coccidiaacidly by destroying all intracellular forms of endogenous development of coccidia-schizonts, micro and macro gametes. The extracellular forms (sporozoites, merozoites and oocysts) are unaffected by toltrazuril. It is active against all pathogenic species of coccidia in birds, toxoplasma, sarcocystis, taylorii. It does not stop the gaining of stable immunity against coccidiosis.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1 ml in 1 L of water (25 ppm with respect to pure toltrazuril) for 2 days or 3 ml in 1L of water (75 ppm) for 8 hours.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Sulfachlorpyrazin-Natrium - 30%

For treatment of coccidiosis. Powder for oral administration with the drinking water.

**Composition in 100g:** Sulfachlorpyrazin-natrium monohydrate: 30g.

**Excipient:** Dextrose monohydrate to 100g.

**Target species:** Pigeons and decorative rabbits.

**Indications:** For treatment of coccidiosis, contagious fever.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g IN 1 L of water for 5 to 7 consecutive days.

**Withdrawal period:** Not applicable.



Package size:

10g 20g 50g 100g 200g 500g 1Kg

Store below 28°C in cool dry place

## Sulfachlorpyrazin-Natrium - 100%

For treatment of coccidiosis. Powder for oral administration with the drinking water.

**Composition in 100g:** Sulfachlorpyrazin-natrium monohydrate: 100g.

**Target species:** Pigeons and decorative rabbits.

**Indications:** Sulfachlorpyrazin has a broad spectrum of activity that covers all types Eimeria causing coccidiosis, as well as Pasteurella multocida, Salmonella gallinarum pulorum.

**Route of administration:** Internally, dissolved in the drinking water.

**Dosage:** 1g in 3 L of water for 3 consecutive days.

*The solution must be prepared daily before use.*

**Withdrawal period:** Not applicable.



Package size:

10ml 20ml 50ml 100ml 500ml 1L

Store below 28°C in cool dry place

## Ectomin 10%

**Insecticide with a residual effect for control of flying insects and creepers. Liquid for water solution.**

**Composition in 100g:** Cypermethrine: 10.0%

**Target species:** Pigeons and decorative animals (guinea pigs, decorative bunnies).

**Indications:** Against flies, mosquitoes, wasps, cockroaches, fleas, ants, ticks, bed bugs, spiders.

**Route of administration:** Spray the walls, wooden and metal panels and windows of the premises. Repeat after 15 days.

**Dosage:** 100ml / 5 l of water.

**Withdrawal period:** Not applicable.



# VETERINARY MEDICINAL PRODUCTS FOR PETS AND LIVESTOCK





**Manufactured by:** „Farma Vet” Ltd., Shumen 9700, Bulgaria



**Worldwide distributor:**

K&K General trading Ltd,  
7, Nish, Str. , Plovdiv 4002, Bulgaria  
phone: + 359 888 262608, +359 899 100537

[www.farmavet.bg](http://www.farmavet.bg)